



CLAIMS

What is claimed is:

1. An amyloid-targeting imaging agent of the formula



- 5 where z is 0 or 1; A_t is an amyloid targeting moiety; A_{lnk} is a linker moiety; and A_{lab} is a labeling moiety.
2. The amyloid-targeting imaging agent of claim 1, where A_t is capable of crossing the blood-brain barrier.
3. The amyloid-targeting imaging agent of claim 1, where A_t is of the formula



10

wherein R^1 and R^2 are each independently a hydrogen atom or a substituted or unsubstituted aliphatic or aryl group; Z and Q^1 are each independently carbonyl ($C=O$), thiocarbonyl ($C=S$), sulfonyl (SO_2), or sulfoxide ($S=O$); k and m are independently 0 or 1, provided that when k is 1, R^1 is not a hydrogen atom and when m is 1, R^2 is not a hydrogen atom; p and s are each independently positive integers selected such that the biodistribution of the targeting moiety for an intended target site is not prevented while maintaining therapeutic activity; T is a linking group; and Y is a group of the formula $-AX$, wherein A is an anionic group at physiological pH, and X is a cationic group.

15


4. The amyloid-targeting imaging agent of claim 3, where R^1 is an alkyl, alkenyl, or aryl group; k is one; Z is a carbonyl group; R^2 is a hydrogen atom or an alkyl group; m is zero; p and s are 1; T is an alkylene group; and Y^1 is SO_3X^2 , where X^2 is H or another cation.

20

5. The amyloid-targeting imaging agent of claim 3, where R^1 and R^2 are alkyl, alkenyl, or aryl, or R^1 and R^2 , taken together, form an alkylene group; k and m are each one; Z and Q^1 are carbonyl groups; p and s are 1; T is an alkylene group; and Y^1 is SO_3X^2 , where X^2 is H or another cation.


25

6. The amyloid-targeting imaging agent of claim 3, where R^1 is an alkyl, alkenyl, or aryl; k and m are zero; R^2 is hydrogen or an alkyl group, p and s are each one; T is an alkylene group; and Y^1 is SO_3X^2 , wherein X^2 is H^+ or another cation.
7. The amyloid-targeting imaging agent of claim 3, where R^1 and R^2 are alkyl, alkenyl, or aryl, or R^1 and R^2 , taken together, form an alkylene group; k and m are zero; p and s are each one; T is an alkylene group; and Y^1 is SO_3X^2 , where X^2 is H^+ or another cation.
8. The amyloid-targeting imaging agent of claim 1, wherein A_i is of formula



wherein Y^- is an anionic group at physiological pH; Q^b is a carrier molecule; X^+ is a cationic group; and n^2 is an integer selected such that the biodistribution of the targeting moiety for the intended target site is not prevented while maintaining activity of the targeting moiety.

9. The amyloid-targeting imaging agent of claim 8, wherein Y is a sulfonate group.
10. The amyloid-targeting imaging agent of claim 8, wherein Y is a sulfate or thiosulfate group.
11. The amyloid-targeting imaging agent of claim 8, wherein Y is a tetrazole group.
12. The amyloid-targeting imaging agent of claim 3, wherein at least one of k or m equals 1.
13. The amyloid-targeting imaging agent of claim 1, where A_i is a peptide of Formula II, an isomer thereof in which the amino acids are of either D or L stereochemistry, a retro or a retro-inverso isomer thereof, or a peptidomimetic thereof:



wherein

- P is selected from the group consisting of peptides which interact with at least one region of an amyloid protein selected from the group consisting of β sheet region, macrophage adherence region, and GAG-binding site region, or $A\beta$ (1-42), fragments or derivatives thereof; said peptide being comprised of natural or unnatural amino acids of either D or L stereochemical configuration;


R' is an N-terminal substituent selected from the group consisting of:

- hydrogen;
- substituted or unsubstituted lower alkyl groups selected from the group consisting of acyclic or cyclic having 1 to 8 carbon atoms;
- aromatic groups;
- heterocyclic groups; and
- acyl groups; and

R'' is a C-terminal substituent selected from the group consisting of hydroxy, alkoxy, aryloxy, unsubstituted or substituted amino groups.

14. The amyloid-targeting imaging agent of claim 13, wherein an amino acid of said peptide of Formula II is a hydrophobic amino acid residue.
15. The amyloid-targeting imaging agent of claim 14, wherein said hydrophobic amino acid residue is a leucine residue.
16. The amyloid-targeting imaging agent of claim 13, wherein said peptide of Formula II has at least two [D] amino acid residues.
17. The amyloid-targeting imaging agent of claim 13, wherein said peptide of Formula II has at least three [D] amino acid residues.
18. The amyloid-targeting imaging agent of claim 13, wherein said peptide of Formula II has one [L] amino acid residue.
19. The amyloid-targeting imaging agent of claim 13, wherein said peptide of Formula II is an all-[D] isomer peptide.
20. The amyloid-targeting imaging agent of claim 1, wherein A_{lnk} is selected from the group consisting of amino, alkylamino, arylamino, oxo, alkoxy, oxoalkyl, aryloxy, oxoaryl, thio, alkylthio, thioalkyl, arylthio, thioaryl, carbonyl, alkylcarbonyl, carbonylalkyl, arylcarbonyl, carbonylaryl, carboxyl, alkylcarboxyl, arylcarboxyl, alkyl, alkylenyl, alkenyl, alkynyl, and aryl groups; glucose; and Phe.
21. The amyloid-targeting imaging agent of claim 1, wherein A_{lab} includes a radionuclide selected from ^{99m}Tc, ⁹⁹Tc, ⁶⁴Cu, ⁶⁷Cu, ⁹⁷Ru, ¹⁰⁹Pd, ¹⁸⁶Re, ¹⁸⁸Re, ¹¹¹In, ^{113m}In, ¹⁵³Gd, ⁹⁰Y, ¹⁵³Sm, ¹⁶⁶Ho, ¹⁹⁸Au, ¹⁹⁹Au, ⁹⁰Sr, ⁸⁹Sr, ¹⁰⁵Rh, ²⁰¹Tl, ⁵¹Cr, ⁶⁷Ga, ⁵⁷Co, ⁶⁰Co, ¹²³I, ¹²⁵I, ¹³¹I or ¹⁸F.

22. The amyloid-targeting imaging agent of claim 1, wherein A_{lab} includes a radionuclide selected from the group consisting of Tc and Re.
23. The amyloid-targeting imaging agent of claim 1, wherein A_{lab} is a metal chelate of a radioactive or paramagnetic metal ion.
- 5 24. The amyloid-targeting imaging agent of claim 1, wherein A_{lab} comprises a chelating ligand of the formula



- where R^{10} is a linear or branched, saturated or unsaturated C_{1-4} alkylene group interrupted by one or two heteroatoms; R^{11} is H or R^{10} , or R^{10} and R^{11} taken together, form a 5- to 8-membered saturated or unsaturated heterocyclic ring optionally substituted with one or more of halogen, hydroxyl, amino, carboxyl, oxo, C_{1-4} alkyl, aryl, or $C(O)R$ groups; R^3 , R^4 , R^5 and R^6 are independently H, carboxyl, C_{1-4} alkyl, an alpha carbon side chain of a D- or L-amino acid other than proline, or $C(O)R$; R^7 and R^8 are independently H, carboxyl, amino, C_{1-4} alkyl, C_{1-4} alkyl; R^9 is H or a sulfur protecting group; and L is hydroxyl, alkoxy, an amino acid residue, or a linking group.
- 10 25. The amyloid-targeting imaging agent of claim 1, wherein A_t is selected from the group consisting of 3-[2-(5-amino-1,2,3,4-tetrahydro isoquinoliny)]-1-propane sulfonic acid hydrochloride, 3[2-(5-bromo-1,2,3,4-tetrahydro isoquinoliny)]-1-propane sulfonic acid, 2-(3-sulfopropyl)-7-amino-1,2,3,4-tetrahydroisoquinoline hydrochloride, 2-(3-sulfopropyl)-7-bromo-1,2,3,4-tetrahydroisoquinoline), Congo Red, 3-(3,4-dihydro-1H-isoquinolin-2-yl)-propane-1-sulfonic acid, 3-(1,3,4,9-tetrahydro-b-carbolin-2-yl)-propane-1-sulfonic acid, 4-(1,3,4,9-tetrahydro-b-carbolin-2-yl)-butane-1-sulfonic acid, 2-amino-3-(3-sulfomethyl-phenyl)-propionic acid, 2-amino-3-(3-sulfomethyl-phenyl)-propionic acid, 2-{4-[(2-amino-4-hydroxy-pteridin-6-ylmethyl)-amino]-benzoylamino}-pentanedioic acid (folic acid), 2,5-dihydroxy-benzene-1,4-disulfonic acid, 2-(4-dimethylamino-phenyl)-3,6-dimethyl-benzothiazol-3-ium; chloride, sodium; 3-(benzothiazol-2-ylsulfanyl)-propane-1-sulfonate, 2,3-dimethyl-benzothiazol-3-ium;
- 15 20 25


iodide, 3-ethyl-2-methyl-benzothiazol-3-ium; iodide, 4-[2-(4-dimethylamino-phenyl)-vinyl]-1-methyl-pyridinium; iodide, 2-[2-(4-dimethylamino-phenyl)-vinyl]-1-ethyl-pyridinium; iodide, and dimethyl-(3-sulfo-propyl)-tetradecyl-ammonium.

26. The amyloid-targeting imaging agent of claim 1, wherein A_i is selected from the group consisting of 2-(3-Sulfo-propyl)-1,2,3,4-tetrahydro-isoquinolin-5-yl-ammonium; chloride; 5-dDiacetyl-amino-2-(3-sulfo-propyl)-isoquinolinium; 5-Nitro-2-(3-sulfo-propyl)-isoquinolinium; 3-(5-bromo-3,4-dihydro-1H-isoquinolin-2-yl)-propane-1-sulfonic acid; 3-(7-nitro-3,4-dihydro-1H-isoquinolin-2-yl)-propane-1-sulfonic acid; 2-(3-sulfo-propyl)-1,2,3,4-tetrahydro-isoquinolin-7-yl-ammonium; chloride; 3-(7-bromo-3,4-dihydro-1H-isoquinolin-2-yl)-propane-1-sulfonic acid; 2-(3-isobutoxysulfonyl-propyl)-5-methyl-1,2,3,4-tetrahydro-isoquinolinium; chloride; 3-(5-iodo-3,4-dihydro-1H-isoquinolin-2-yl)-propane-1-sulfonic acid; 2-(3-sulfo-propyl)-9H-b-carbolin-2-ium; 2-(2-methoxycarbonyl-ethyl)-1,2,3,4-tetrahydro-isoquinolinium; chloride; 2-(3-sulfo-propyl)-1,2,3,4-tetrahydro-isoquinolin-6-yl-ammonium; chloride; 3-(6-bromo-3,4-dihydro-1H-isoquinolin-2-yl)-propane-1-sulfonic acid; 2-(3-sulfo-propyl)-1,2,3,4-tetrahydro-isoquinoline-6-carboxylic acid methyl ester; 2-(3-sulfo-propyl)-1,2,3,4-tetrahydro-isoquinoline-5-carboxylic acid methyl ester; 2-(3-sulfo-propyl)-1,2,3,4-tetrahydro-isoquinoline-7-carboxylic acid methyl ester; 3-(6-bromo-1,3,4,9-tetrahydro-b-carbolin-2-yl)-propane-1-sulfonic acid; 3-(6-amino-1,3,4,9-tetrahydro-b-carbolin-2-yl)-propane-1-sulfonic acid; 2-(3-sulfo-propyl)-2,3,4,9-tetrahydro-1H-b-carboline-6-carboxylic acid methyl ester; 4-(6-bromo-1,3,4,9-tetrahydro-b-carbolin-2-yl)-butane-1-sulfonic acid; 4-(6-amino-1,3,4,9-tetrahydro-b-carbolin-2-yl)-butane-1-sulfonic acid; and 2-(4-sulfo-butyl)-2,3,4,9-tetrahydro-1H-b-carboline-6-carboxylic acid methyl ester.
27. The amyloid-targeting imaging agent of claim 1, wherein A_i is selected from the group consisting of 3-phenylamino-1-propanesulfonic acid sodium salt, 3-(4-pyridylamino)]-1-propanesulfonic acid, 3-(benzylamino)-1-propanesulfonic acid, diethylphosphonoacetic acid, phosphonoformic acid, trisodium salt, 3-benzoylamino-propanesulfonic acid, 2-deoxy-2-(3-sulfopropyl)amino-d-glucose, 1-phenyl-2,3,-dimethyl-4-methylamino-pyrazolon-5-N-methylsulfonic acid, 3-[(3,4-dimethyl-1-adamantyl)-amino]-1-propanesulfonic acid, 3-(2-hydroxyethyl)amino-1-propanesulfonic acid, 3-(3-hydroxy-1-propyl)amino-1-propanesulfonic acid, (-)-3-[(R)-2-hydroxy-1-propyl]amino-1-propanesulfonic acid, 3-[(d,l)-2-hydroxy-1-propyl]-1-

- propanesulfonic acid, 3-(4-hydroxy-1-butyl)amino-1-propanesulfonic acid, 3-(5-hydroxy-1-pentyl)amino-1-propanesulfonic acid, 3-(6-hydroxy-1-hexyl)amino-1-propanesulfonic acid, 3-(4-hydroxyphenyl)amino-1-propanesulfonic acid, (+)-3-[(S)-2-hydroxy-1-propyl]amino-1-propanesulfonic acid, (+)-3-[(S)-1-hydroxy-2-propyl]amino-1-propanesulfonic acid, (-)-3-[(R)-1-hydroxy-2-propyl]amino-1-propanesulfonic acid, (+)-3-[(S)-1-hydroxy-2-butyl]amino-1-propanesulfonic acid, (-)-3-[(R)-1-hydroxy-2-butyl]amino-1-propanesulfonic acid, 3-[(dl)-5-hydroxy-2-pentyl]amino-1-propanesulfonic acid, 3-[(dl)-6-hydroxy-2-hexyl]amino-1-propanesulfonic acid, 3-amylamino-1-propanesulfonic acid, 3-hexylamino-1-propanesulfonic acid, 3-heptylamino-1-propanesulfonic acid, 3-octylamino-1-propanesulfonic acid, 3-nonylamino-1-propanesulfonic acid, 3-decylamino-1-propanesulfonic acid, 3-undecylamino-1-propanesulfonic acid, 3-dodecylamino-1-propanesulfonic acid, 3-tridecylamino-1-propanesulfonic acid, 3-tetradecylamino-1-propanesulfonic acid, 3-hexadecylamino-1-propanesulfonic acid, 3-octadecylamino-1-propanesulfonic acid, dimethyl(3-sulfopropyl)-tetradecylammonium hydroxide, inner salt, 2-(3-Sulfobutyl)-1,2,3,4-tetrahydro-9H-pyrido [3,4-b] indole, sodium, 3-[1,2,3,4-tetrahydro-9H-pyrido (3,4-b)indolyl]-1-propanesulfonic acid, 2,5-dihydroxy-1,4-benzenedisulfonic acid, H-dLys-dLeu-dVal-dPhe-dPhe-dAla-OH (SEQ ID NO. 28), Thioflavin T, Folic acid dihydrate, 3-(2-benzothiazolylthio)-1-propanesulfonic acid, 2,3-dimethylbenzothiazolium, 3-ethyl-2-methylbenzothiazolium, trans-4-[4-(dimethylamino)styryl]-1-methylpyridinium, 2-[4-(dimethylamino)styryl]-1-ethylpyridinium, and dimethyl(3-sulfopropyl) tetradecylammonium hydroxide inner salt.
28. The amyloid-targeting imaging agent of claim 1, wherein A₁ is selected from the group consisting of 3-acetylaminopropanesulfonic acid, 3-benzoylamino-1-propanesulfonic acid sodium salt, and 2-acrylamido-2-methyl-1-propanesulfonic acid.
29. The amyloid-targeting imaging agent of claim 1, wherein A₁ is selected from the group consisting of 3-phthalimido-1-propanesulfonic acid, N-(3-sulfopropyl)saccharin and 4-phthalimido-1-butanedisulfonic acid.
30. The amyloid-targeting imaging agent of claim 1, wherein A₁ is selected from the group consisting of 3-dimethylamino-1-propanesulfonic acid, 4-(1-piperidinyl)-1-butanedisulfonic acid, 3-[1-(1,2,3,6-tetrahydropyridyl)]-1-propanesulfonic acid, 3-[2-

- (1,2,3,4-tetrahydroisoquinoliny)]-1-propanesulfonic acid, 3-[2-(6,7-dimethoxy-1,2,3,4-tetrahydroisoquinoliny)]-1-propanesulfonic acid, 3-[1-(1,2,3,4-tetrahydroisoquinoliny)]-1-propanesulfonic acid, 2-(3-sulfopropyl)-1,2,3,4-tetrahydro-9H-pyrido[3,4-b]indole, sodium salt, 3-(1-indoliny)]-1-propanesulfonic acid, 3-[2-(6-methoxy-1,2,3,4-tetrahydroisoquinoliny)]-1-propanesulfonic acid, 3-(2-isoindoliny)]-1-propanesulfonic acid, 3-(4-benzyl-1-piperidiny)]-1-propanesulfonic acid, 1-(3-sulfopropyl)-(S)-nicotinium hydroxide inner salt, 3-[2-(1,2,3,4,5,6,7,8-octahydroisoquinoliny)]-1-propanesulfonic acid, Thiazol Yellow G, 3-sulfolmethylphenylalanine, Chicago Sky Blue 6B, 4-[2-(1,2,3,4-tetrahydroisoquinoliny)]-1-butan-sulfonic acid, and 3-sulfomethyl-L-phenylalanine.


31. The amyloid-targeting imaging agent of claim 1, where A_t is of the formula



where

- R^1 is an alkyl, alkenyl, hydroxyalkyl, or a single-ring aromatic group;
- R^2 is a alkyl, alkenyl, hydroxyalkyl, a single-ring aromatic group, or a hydrogen atom, or R^1 and R^2 , taken together with the nitrogen to which they are attached, form a heterocyclic group which is a fused ring structure;
- T is an alkylene group;
- Y is SO_3X , and X is a cationic group.


32. The amyloid-targeting imaging agent of claim 1, where A_t is of the formula



where

- R^1 is a C_5 - C_{18} alkyl, hydroxyalkyl or single-ring aromatic group;
- R^2 is a hydrogen atom or an alkyl group;
- T is an alkylene group;
- Y is SO_3X , and X is a cationic group.


33. The amyloid-targeting imaging agent of claim 1, where A_t is of the formula



where

- R^1 is an alkyl, an alkenyl, or an aromatic group;
- R^2 is a hydrogen atom, an alkyl group, or an aromatic group, or R^1 and R^2 , taken together, form a heterocyclic group which is a fused ring structure;
- Z and Q are each independently a carbonyl (C=O), thiocarbonyl (C=S), sulfonyl (SO₂), or sulfoxide (S=O) group;
- k is 1 and m is 0 or 1;
- p and s are each 1;
- T is an alkylene group;
- Y is SO₃X, and X is a cationic group.

34. The amyloid-targeting imaging agent of claim 1, where A_t is of the formula



where

- R^1 and R^2 are alkyl, alkenyl, or single-ring aromatic groups, or R^1 and R^2 , taken together with the nitrogen to which they are attached, form a heterocyclic group which is a fused ring structure;
- T is an alkylene group;
- Y is SO₃X, and X is a cationic group.


35. The amyloid-targeting imaging agent of claim 33, wherein said A_t is selected from the group consisting of 3-acetylamino-1-propanesulfonic acid, 3-benzoylamino-1-propanesulfonic acid, and 2-acrylamido-2-methyl-1-propanesulfonic acid.

36. The amyloid-targeting imaging agent of claim 33, wherein said A_t is selected from the group consisting of 3-phthalimido-1-propanesulfonic acid, N-(3-sulfopropyl)saccharin and 4-phthalimido-1-butanefulfonic acid.

37. The amyloid-targeting imaging agent of claim 32, wherein said A₁ is selected from the group consisting of 3-phenylamino-1-propanesulfonic acid, 3-(4-pyridylamino)-1-propanesulfonic acid, 3-(benzylamino)-1-propanesulfonic acid, 2-deoxy-2-(3-sulfopropyl)amino-D-glucose, 3-[(3,4-dimethyl-1-adamantyl)-amino]-1-propanesulfonic acid, 3-[(3,5-dimethyl-1-adamantyl)-amino]-1-propanesulfonic acid, 2-deoxy-2-(3-sulfopropyl)amino-D-glucose, 3-[(3,5-dimethyl-1-adamantyl)-amino]-1-propanesulfonic acid, 3-(2-hydroxyethyl)amino-1-propanesulfonic acid, 3-(3-hydroxy-1-propyl)amino-1-propanesulfonic acid, (-)-3-[(R)-2-hydroxy-1-propyl]amino-1-propanesulfonic acid, 3-[(d,l)-1-hydroxy-2-propyl]amino-1-propanesulfonic acid, 3-(4-hydroxy-1-butyl)amino-1-propanesulfonic acid, 3-(5-hydroxy-1-pentyl)amino-1-propanesulfonic acid, 3-(6-hydroxy-1-hexyl)amino-1-propanesulfonic acid, 3-(4-hydroxyphenyl)amino-1-propanesulfonic acid, (+)-3-[(S)-2-hydroxy-1-propyl]amino-1-propanesulfonic acid, (+)-3-[(S)-1-hydroxy-2-propyl]amino-1-propanesulfonic acid, (-)-3-[(R)-1-hydroxy-2-propyl]amino-1-propanesulfonic acid, (+)-3-[(S)-1-hydroxy-2-butyl]amino-1-propanesulfonic acid, (-)-3-[(R)-1-hydroxy-2-butyl]amino-1-propanesulfonic acid, 3-[(dl)-5-hydroxy-2-pentyl]amino-1-propanesulfonic acid, 3-[(dl)-6-hydroxy-2-hexyl]amino-1-propanesulfonic acid, 3-(1-hydroxymethyl-1-cyclopentyl)amino-1-propanesulfonic acid, 3-amylamino-1-propanesulfonic acid, 3-hexylamino-1-propanesulfonic acid, 3-heptylamino-1-propanesulfonic acid, 3-octylamino-1-propanesulfonic acid, 3-nonylamino-1-propanesulfonic acid, 3-decylamino-1-propanesulfonic acid, 3-undecylamino-1-propanesulfonic acid, 3-dodecylamino-1-propanesulfonic acid, 3-tridecylamino-1-propanesulfonic acid, 3-tetradecylamino-1-propanesulfonic acid, 3-hexadecylamino-1-propanesulfonic acid, and 3-octadecylamino-1-propanesulfonic acid.
38. The amyloid-targeting imaging agent of claim 34, wherein said A₁ is selected from the group consisting of 1-phenyl-2,3-dimethyl-4-methylamino-pyrazolon-5-N-methylsulfonic acid; 3-dimethylamino-1-propanesulfonic acid, 3-[2-(1,2,3,4-tetrahydroisoquinoliny)]-1-propanesulfonic acid, 3-[2-(6,7-dimethoxy-1,2,3,4-tetrahydroisoquinoliny)]-1-propanesulfonic acid, 3-[1-(1,2,3,4-tetrahydroquinoliny)]-1-propanesulfonic acid, 2-(3-sulfopropyl)-1,2,3,4-tetrahydro-9H-pyrido[3,4-b]indole, 3-(1-indoliny)]-1-propanesulfonic acid, 3-[2-(6-methoxy-1,2,3,4-tetrahydroisoquinoliny)]-1-propanesulfonic acid, 3-(2-isoindoliny)]-1-propanesulfonic

acid, 3-[2-(1,2,3,4,5,6,7,8-octahydroisoquinoliny)]-1-propanesulfonic acid, and 4-[2-(1,2,3,4-tetrahydroisoquinoliny)]-1-butanefulfonic acid.

39. The amyloid-targeting imaging agent of claim 1, where said A_t is selected from the group consisting of 4-phenyl-1-(3'-sulfopropyl)-1,2,3,6-tetrahydropyridine; 2-(3-sulfobutyl)-1,2,3,4-tetrahydro-9H-pyrido [3,4-b] indole; 3-(4-benzyl-1-piperidiny)-1-propanesulfonic acid, 3-sulfonylmethylphenylalanine, 4-(1-piperidiny)-1-butanefulfonic acid, cyclohexylsulfamic acid; 1-(3-sulfopropyl)-(S)-nicotinium hydroxide inner salt, 3-[1-(1,2,3,6-tetrahydropyridyl)]-1-propanesulfonic acid, 3-sulfomethyl-D,L-phenylalanine and 3-sulfomethyl-L-phenylalanine.
40. A method for diagnosing an amyloid-related condition in a patient, comprising administering an amyloid-targeting imaging agent to a patient, and imaging said amyloid-targeting imaging agent in said patient to determine the presence of amyloid in said patient, such that the presence or absence of an amyloid-related condition in said patient is determined.
41. The method of claim 40, wherein amyloid-targeting imaging agent is of the formula




where z is 0 or 1; A_t is an amyloid targeting moiety; A_{lnk} is a linker moiety; and A_{lab} is a labeling moiety.

42. The method of claim 41, wherein said amyloid-related condition is selected from the group consisting of Creutzfeld-Jakob Disease (CJD), Kuru, transmissible cerebral amyloidoses (also known as transmissible virus dementias), familial CJD, scrapie, transmissible mink encephalopathy, bovine spongiform encephalopathy (BSE), inflammation-associated amyloid, type II diabetes, primary amyloidosis, feline spongiform encephalopathy, non-transmissible cerebral amyloidosis (e.g., Alzheimer's disease), prion-mediated diseases, dialysis-related amyloidosis, light chain-related amyloidosis, cerebral amyloid angiopathy, and Alzheimer's disease.
43. A kit for preparing a radiopharmaceutical preparation, said kit comprising:
- an amyloid-targeting imaging agent of claim 1;
 - a reducing agent;

- a buffering agent;
- a transchelating agent, and
- instructions for the preparation and use of the radiopharmaceutical in the imaging of amyloid or an amyloid-related condition.

5 44. The kit of claim 43, wherein A_1 is of formula



wherein Y^- is an anionic group at physiological pH; Q^b is a carrier molecule; X^+ is a cationic group; and n^2 is an integer selected such that the biodistribution of the targeting moiety for the intended target site is not prevented while maintaining activity of the targeting moiety.

10

45. The kit of claim 44, wherein Y is a sulfonate group.

46. The kit of claim 44, wherein Y is a sulfate or thiosulfate group.

47. The kit of claim 44, wherein Y is a tetrazole group.

15

48. A method for imaging amyloid deposition in a patient, comprising administering an amyloid-targeting imaging agent to a patient, and imaging said amyloid-targeting imaging agent in said patient to determine the presence of amyloid in said patient.

49. The amyloid-targeted imaging agent of claim 13 wherein said peptide is selected from the group consisting of SEQ ID NOS 1-28.

20

50. The amyloid-targeted imaging agent of claim 49 wherein said peptide is modified by substitution of one amino acid by a different amino acid or by deletion of one amino acid.


51. A method of diagnostic medical imaging of an amyloid-associated disease comprising the steps of administering to a patient a pharmaceutical composition according to claim 1 and then imaging said patient.

25

52. The method of diagnostic medical imaging according to claim 51 wherein A_{lab} of said pharmaceutical composition is a radiopharmaceutical.

53. The method of diagnostic medical imaging according to claim 51 wherein A_{lab} of said pharmaceutical composition is a metal chelate.


54. The method of diagnostic medical imaging according to claim 53 wherein said metal chelate is gadolinium-DTPA, gadolinium-DOTA, or gadolinium-DO3A.
55. The method of diagnostic medical imaging according to claim 53 wherein said metal chelate is a chelate of ^{99m}Tc or ^{111}In .
- 5 56. The method of diagnostic medical imaging according to claim 51 wherein said imaging step is ultrasound imaging.
57. The method of claim 48, wherein said imaging step is radionuclide imaging.
58. The method of claim 57, wherein said imaging step is SPECT imaging.
59. The method of claim 48, wherein said imaging step is magnetic resonance imaging.
- 10 60. The method of claim 48, wherein said imaging step is ultrasound imaging.
61. The method of claim 48, wherein said imaging step is X-ray imaging.
62. The method of claim 48, wherein said imaging step is fluorescence imaging.
63. The method of claim 48, wherein said amyloid-targeting imaging agent has the formula



where z is 0 or 1; A_t is an amyloid targeting moiety; A_{lnk} is a linker moiety; and A_{lab} is a labeling moiety.

64. The method of claim 63, wherein A_{lnk} is selected from the group consisting of amino, 20 alkylamino, arylamino, oxo, alkoxy, oxoalkyl, aryloxy, oxoaryl, thio, alkylthio, thioalkyl, arylthio, thioaryl, carbonyl, alkylcarbonyl, carbonylalkyl, arylcarbonyl, carbonylaryl, carboxyl, alkylcarboxyl, arylcarboxyl, alkyl, alkylenyl, alkenyl, alkynyl, and aryl groups; glucose; and Phe.
65. The method of claim 63, wherein A_{lab} includes a radionuclide selected from ^{99m}Tc , ^{99}Tc , 25 ^{64}Cu , ^{67}Cu , ^{97}Ru , ^{109}Pd , ^{186}Re , ^{188}Re , ^{111}In , ^{113m}In , ^{153}Gd , ^{90}Y , ^{153}Sm , ^{166}Ho , ^{198}Au , ^{199}Au , ^{90}Sr , ^{89}Sr , ^{105}Rh , ^{201}Tl , ^{51}Cr , ^{67}Ga , ^{57}Co , ^{60}Co , ^{123}I , ^{125}I , ^{131}I , or ^{18}F .
66. The method of claim 63, wherein A_{lab} includes a radionuclide selected from the group consisting of Tc and Re.

67. The method of claim 63, wherein A_{lab} is a metal chelate of a radioactive or paramagnetic metal ion.
68. The method of claim 63, wherein A_{lab} comprises a chelating ligand of the formula




5

where R^{10} is a linear or branched, saturated or unsaturated C_{1-4} alkylene group interrupted by one or two heteroatoms; R^{11} is H or R^{10} , or R^{10} and R^{11} taken together, form a 5- to 8-membered saturated or unsaturated heterocyclic ring optionally substituted with one or more of halogen, hydroxyl, amino, carboxyl, oxo, C_{1-4} alkyl, aryl, or $C(O)R$ groups; R^3 , R^4 , R^5 and R^6 are independently H, carboxyl, C_{1-4} alkyl, an alpha carbon side chain of a D- or L-amino acid other than proline, or $C(O)R$; R^7 and R^8 are independently H, carboxyl, amino, C_{1-4} alkyl, C_{1-4} alkyl; R^9 is H or a sulfur protecting group; and L is hydroxyl, alkoxy, an amino acid residue, or a linking group.

10

69. The method of claim 63, where A_i is capable of crossing the blood-brain barrier.
70. The method of claim 63, where A_i is of the formula




20

wherein R^1 and R^2 are each independently a hydrogen atom or a substituted or unsubstituted aliphatic or aryl group; Z and Q^1 are each independently carbonyl ($C=O$), thiocarbonyl ($C=S$), sulfonyl (SO_2), or sulfoxide ($S=O$); k and m are independently 0 or 1, provided that when k is 1, R^1 is not a hydrogen atom and when m is 1, R^2 is not a hydrogen atom; p and s are each independently positive integers selected such that the

biodistribution of the targeting moiety for an intended target site is not prevented while maintaining therapeutic activity; T is a linking group; and Y is a group of the formula -AX, wherein A is an anionic group at physiological pH, and X is a cationic group.

71. The method of claim 70, where R^1 is an alkyl, alkenyl, or aryl group; k is one; Z is a carbonyl group; R^2 is a hydrogen atom or an alkyl group; m is zero; p and s are 1; T is an alkylene group; and Y^1 is SO_3X^2 , where X^2 is H or another cation.
72. The method of claim 70, where R^1 and R^2 are alkyl, alkenyl, or aryl, or R^1 and R^2 , taken together, form an alkylene group; k and m are each one; Z and Q^1 are carbonyl groups; p and s are 1; T is an alkylene group; and Y^1 is SO_3X^2 , where X^2 is H or another cation.
73. The method of claim 70, where R^1 is an alkyl, alkenyl, or aryl; k and m are zero; R^2 is hydrogen or an alkyl group, p and s are each one; T is an alkylene group; and Y^1 is SO_3X^2 , wherein X^2 is H^+ or another cation.
74. The method of claim 70, where R^1 and R^2 are alkyl, alkenyl, or aryl, or R^1 and R^2 , taken together, form an alkylene group; k and m are zero; p and s are each one; T is an alkylene group; and Y^1 is SO_3X^2 , where X^2 is H^+ or another cation.
75. The method of claim 63, wherein A_t is of formula




- wherein Y^- is an anionic group at physiological pH; Q^b is a carrier molecule; X^+ is a cationic group; and n^2 is an integer selected such that the biodistribution of the targeting moiety for the intended target site is not prevented while maintaining activity of the targeting moiety.
76. The method of claim 75, wherein Y is a sulfonate group.
77. The method of claim 75, wherein Y is a sulfate or thiosulfate group.
78. The method of claim 75, wherein Y is a tetrazole group.
79. The method of claim 70, wherein at least one of k or m equals 1.

80. The method of claim 63, wherein A_i is selected from the group consisting of 3-[2-(5-amino-1,2,3,4-tetrahydro isoquinolinyl)]-1-propane sulfonic acid hydrochloride, 3[2-(5-bromo-1,2,3,4-tetrahydro isoquinolinyl)]-1-propane sulfonic acid, 2-(3-sulfopropyl)-7-amino-1,2,3,4-tetrahydroisoquinoline hydrochloride, 2-(3-sulfopropyl)-7-bromo-1,2,3,4-tetrahydroisoquinoline), Congo Red, 3-(3,4-dihydro-1H-isoquinolin-2-yl)-propane-1-sulfonic acid, 3-(1,3,4,9-tetrahydro-b-carbolin-2-yl)-propane-1-sulfonic acid, 4-(1,3,4,9-tetrahydro-b-carbolin-2-yl)-butane-1-sulfonic acid, 2-amino-3-(3-sulfomethyl-phenyl)-propionic acid, 2-amino-3-(3-sulfomethyl-phenyl)-propionic acid, 2-{4-[(2-amino-4-hydroxy-pteridin-6-ylmethyl)-amino]-benzoylamino}-pentanedioic acid, 2,5-dihydroxy-benzene-1,4-disulfonic acid, 2-(4-dimethylamino-phenyl)-3,6-dimethyl-benzothiazol-3-ium; chloride, sodium; 3-(benzothiazol-2-ylsulfanyl)-propane-1-sulfonate, 2,3-dimethyl-benzothiazol-3-ium; iodide, 3-ethyl-2-methyl-benzothiazol-3-ium; iodide, 4-[2-(4-dimethylamino-phenyl)-vinyl]-1-methyl-pyridinium; iodide, 2-[2-(4-dimethylamino-phenyl)-vinyl]-1-ethyl-pyridinium; iodide, and dimethyl-(3-sulfo-propyl)-tetradecyl-ammonium.
81. The method of claim 63, wherein A_i is selected from the group consisting of 2-(3-Sulfo-propyl)-1,2,3,4-tetrahydro-isoquinolin-5-yl-ammonium; chloride; 5-diacetylamino-2-(3-sulfo-propyl)-isoquinolinium; 5-Nitro-2-(3-sulfo-propyl)-isoquinolinium; 3-(5-bromo-3,4-dihydro-1H-isoquinolin-2-yl)-propane-1-sulfonic acid; 3-(7-nitro-3,4-dihydro-1H-isoquinolin-2-yl)-propane-1-sulfonic acid; 2-(3-sulfo-propyl)-1,2,3,4-tetrahydro-isoquinolin-7-yl-ammonium; chloride; 3-(7-bromo-3,4-dihydro-1H-isoquinolin-2-yl)-propane-1-sulfonic acid; 2-(3-isobutoxysulfonyl-propyl)-5-methyl-1,2,3,4-tetrahydro-isoquinolinium; chloride; 3-(5-iodo-3,4-dihydro-1H-isoquinolin-2-yl)-propane-1-sulfonic acid; 2-(3-sulfo-propyl)-9H-b-carbolin-2-ium; 2-(2-methoxycarbonyl-ethyl)-1,2,3,4-tetrahydro-isoquinolinium; chloride; 2-(3-sulfo-propyl)-1,2,3,4-tetrahydro-isoquinolin-6-yl-ammonium; chloride; 3-(6-bromo-3,4-dihydro-1H-isoquinolin-2-yl)-propane-1-sulfonic acid; 2-(3-sulfo-propyl)-1,2,3,4-tetrahydro-isoquinoline-6-carboxylic acid methyl ester; 2-(3-sulfo-propyl)-1,2,3,4-tetrahydro-isoquinoline-5-carboxylic acid methyl ester; 2-(3-sulfo-propyl)-1,2,3,4-tetrahydro-isoquinoline-7-carboxylic acid methyl ester; 3-(6-bromo-1,3,4,9-tetrahydro-b-carbolin-2-yl)-propane-1-sulfonic acid; 3-(6-amino-1,3,4,9-tetrahydro-b-carbolin-2-yl)-propane-1-sulfonic acid; 2-(3-sulfo-propyl)-2,3,4,9-tetrahydro-1H-b-carboline-6-carboxylic acid methyl ester; 4-(6-bromo-1,3,4,9-tetrahydro-b-carbolin-2-yl)-butane-1-sulfonic acid; 4-(6-amino-

1,3,4,9-tetrahydro-b-carbolin-2-yl)-butane-1-sulfonic acid; and 2-(4-sulfo-butyl)-2,3,4,9-tetrahydro-1H-b-carboline-6-carboxylic acid methyl ester.

82. The method of claim 63, wherein A_i is selected from the group consisting of 3-phenylamino-1-propanesulfonic acid sodium salt, 3-(4-pyridylamino)]-1-
5 propanesulfonic acid, 3-(benzylamino)-1-propanesulfonic acid, diethylphosphonoacetic acid, phosphonoformic acid, trisodium salt, 3-benzoylaminopropanesulfonic acid, 2-deoxy-2-(3-sulfopropyl)amino-d-glucose, 1-phenyl-2,3,-dimethyl-4-methylamino-pyrazolon-5-N-methylsulfonic acid, 3-[-(3,4-dimethyl-1-adamantyl)-amino]-1-propanesulfonic acid, 3-(2-hydroxyethyl)amino-1-propanesulfonic acid, 3-(3-hydroxy-
10 1-propyl)amino-1-propanesulfonic acid, (-)-3-[(R)-2-hydroxy-1-propyl]amino-1-propanesulfonic acid, 3-[(d,l)-2-hydroxy-1-propyl]-1-propanesulfonic acid, 3-(4-hydroxy-1-butyl)amino-1-propanesulfonic acid, 3-(5-hydroxy-1-pentyl)amino-1-propanesulfonic acid, 3-(6-hydroxy-1-hexyl)amino-1-propanesulfonic acid, 3-(4-hydroxyphenyl)amino-1-propanesulfonic acid, (+)-3-[(S)-2-hydroxy-1-propyl]amino-1-propanesulfonic acid, (+)-3-[(S)-1-hydroxy-2-propyl]amino-1-propanesulfonic acid, (-)-
15 3-[(R)-1-hydroxy-2-propyl]amino-1-propanesulfonic acid, (+)-3-[(S)-1-hydroxy-2-butyl]amino-1-propanesulfonic acid, (-)-3-[(R)-1-hydroxy-2-butyl]amino-1-propanesulfonic acid, 3-[(dl)-5-hydroxy-2-pentyl]amino-1-propanesulfonic acid, 3-[(dl)-6-hydroxy-2-hexyl]amino-1-propanesulfonic acid, 3-amylamino-1-propanesulfonic acid, 3-hexylamino-1-propanesulfonic acid, 3-heptylamino-1-propanesulfonic acid, 3-octylamino-1-propanesulfonic acid, 3-nonylamino-1-propanesulfonic acid, 3-decylamino-1-propanesulfonic acid, 3-undecylamino-1-propanesulfonic acid, 3-dodecylamino-1-propanesulfonic acid, 3-tridecylamino-1-propanesulfonic acid, 3-tetradecylamino-1-propanesulfonic acid, 3-hexadecylamino-1-propanesulfonic acid, 3-octadecylamino-1-propanesulfonic acid, dimethyl(3-sulfopropyl)-tetradecylammonium hydroxide, inner salt, 2-(3-Sulfobutyl)-1,2,3,4-tetrahydro-9H-pyrido [3,4-b] indole, sodium, 3-[1,2,3,4-tetrahydro-9H-pyrido (3,4-b)indolyl]-1-propanesulfonic acid, 2,5-dihydroxy-1,4-benzenedisulfonic acid, Thioflavin T, Folic acid dihydrate, 3-(2-benzothiazolylthio)-1-propanesulfonic acid,
20 2,3-dimethylbenzothiazolium, 3-ethyl-2-methylbenzothiazolium, trans-4-[4-(dimethylamino)styryl]-1-methylpyridinium, 2-[4-(dimethylamino)styryl]-1-ethylpyridinium, and dimethyl(3-sulfopropyl) tetradecylammonium hydroxide inner salt.


83. The method of claim 63, wherein A_1 is selected from the group consisting of 3-acetylamino propanesulfonic acid, 3-benzoylamino-1-propanesulfonic acid sodium salt, and 2-acrylamido-2-methyl-1-propanesulfonic acid.
- 5 84. The method of claim 63, wherein A_1 is selected from the group consisting of 3-phthalimido-1-propanesulfonic acid, N-(3-sulfopropyl)saccharin and 4-phthalimido-1-butanefulfonic acid.
- 10 85. The method of claim 63, wherein A_1 is selected from the group consisting of 3-dimethylamino-1-propanesulfonic acid, 4-(1-piperidinyl)-1-butanefulfonic acid, 3-[1-(1,2,3,6-tetrahydropyridyl)]-1-propanesulfonic acid, 3-[2-(1,2,3,4-tetrahydroisoquinolyl)]-1-propanesulfonic acid, 3-[2-(6,7-dimethoxy-1,2,3,4-tetrahydroisoquinolyl)]-1-propanesulfonic acid, 3-[1-(1,2,3,4-tetrahydroquinolyl)]-1-propanesulfonic acid, 2-(3-sulfopropyl)-1,2,3,4-tetrahydro-9H-pyrido[3,4-b]indole, sodium salt, 3-(1-indolyl)-1-propanesulfonic acid, 3-[2-(6-methoxy-1,2,3,4-tetrahydroisoquinolyl)]-1-propanesulfonic acid, 3-(2-isoindolyl)-1-propanesulfonic acid, 3-(4-benzyl-1-piperidinyl)-1-propanesulfonic acid, 1-(3-sulfopropyl)-(S)-nicotinium hydroxide inner salt, 3-[2-(1,2,3,4,5,6,7,8-octahydroisoquinolyl)]-1-propanesulfonic acid, Thiazol Yellow G, 3-sulfomethylphenylalanine, Chicago Sky Blue 6B, 4-[2-(1,2,3,4-tetrahydroisoquinolyl)]-1-butanefulfonic acid, and 3-sulfomethyl-L-phenylalanine.
- 15 20 86. The method of claim 63, where A_1 is of the formula



where

- 25 R^1 is an alkyl, alkenyl, hydroxyalkyl, or a single-ring aromatic group;
 R^2 is a alkyl, alkenyl, hydroxyalkyl, a single-ring aromatic group, or a hydrogen atom, or R^1 and R^2 , taken together with the nitrogen to which they are attached, form a heterocyclic group which is a fused ring structure;
T is an alkylene group;
30 Y is SO_3X , and X is a cationic group.

87. The method of claim 63, where A_i is of the formula



5 where


R^1 is a C_5 - C_{18} alkyl, hydroxyalkyl or single-ring aromatic group;

R^2 is a hydrogen atom or an alkyl group;

T is an alkylene group;

Y is SO_3X , and X is a cationic group.

10 88. The method of claim 63, where A_i is of the formula



where

15 R^1 is an alkyl, an alkenyl, or an aromatic group;

R^2 is a hydrogen atom, an alkyl group, or an aromatic group, or R^1 and R^2 , taken together, form a heterocyclic group which is a fused ring structure;

Z and Q are each independently a carbonyl ($C=O$), thiocarbonyl ($C=S$), sulfonyl (SO_2), or sulfoxide ($S=O$) group;

20 k is 1 and m is 0 or 1;


p and s are each 1;

T is an alkylene group;

Y is SO_3X , and X is a cationic group.

89. The method of claim 63, where A_i is of the formula

25



where


R^1 and R^2 are alkyl, alkenyl, or single-ring aromatic groups, or R^1 and R^2 , taken together with the nitrogen to which they are attached, form a heterocyclic group which is a fused ring structure;

T is an alkylene group;

Y is SO_3X , and X is a cationic group.

90. The method of claim 88, wherein said A_i is selected from the group consisting of 3-acetylamino-1-propanesulfonic acid, 3-benzoylamino-1-propanesulfonic acid, and 2-acrylamido-2-methyl-1-propanesulfonic acid.
91. The method of claim 88, wherein said A_i is selected from the group consisting of 3-phthalimido-1-propanesulfonic acid, N-(3-sulfopropyl)saccharin and 4-phthalimido-1-butanefulfonic acid.
92. The method of claim 87, wherein said A_i is selected from the group consisting of 3-phenylamino-1-propanesulfonic acid, 3-(4-pyridylamino)-1-propanesulfonic acid, 3-(benzylamino)-1-propanesulfonic acid, 2-deoxy-2-(3-sulfopropyl)amino-D-glucose, 3-[(3,4-dimethyl-1-adamantyl)-amino]-1-propanesulfonic acid, 3-[(3,5-dimethyl-1-adamantyl)-amino]-1-propanesulfonic acid, 3-(2-hydroxyethyl)amino-1-propanesulfonic acid, 3-(3-hydroxy-1-propyl)amino-1-propanesulfonic acid, (-)-3-[(R)-2-hydroxy-1-propyl]amino-1-propanesulfonic acid, 3-[(d,l)-1-hydroxy-2-propyl]amino-1-propanesulfonic acid, 3-(4-hydroxy-1-butyl)amino-1-propanesulfonic acid, 3-(5-hydroxy-1-pentyl)amino-1-propanesulfonic acid, 3-(6-hydroxy-1-hexyl)amino-1-propanesulfonic acid, 3-(4-hydroxyphenyl)amino-1-propanesulfonic acid, (+)-3-[(S)-2-hydroxy-1-propyl]amino-1-propanesulfonic acid, (+)-3-[(S)-1-hydroxy-2-propyl]amino-1-propanesulfonic acid, (-)-3-[(R)-1-hydroxy-2-propyl]amino-1-propanesulfonic acid, (+)-3-[(S)-1-hydroxy-2-butyl]amino-1-propanesulfonic acid, (-)-3-[(R)-1-hydroxy-2-butyl]amino-1-propanesulfonic acid, 3-[(dl)-5-hydroxy-2-pentyl]amino-1-propanesulfonic acid, 3-[(dl)-6-hydroxy-2-hexyl]amino-1-propanesulfonic acid, 3-(1-hydroxymethyl-1-cyclopentyl)amino-1-propanesulfonic acid, 3-amylamino-1-

- propanesulfonic acid, 3-hexylamino-1-propanesulfonic acid, 3-heptylamino-1-propanesulfonic acid, 3-octylamino-1-propanesulfonic acid, 3-nonylamino-1-propanesulfonic acid, 3-decylamino-1-propanesulfonic acid, 3-undecylamino-1-propanesulfonic acid, 3-dodecylamino-1-propanesulfonic acid, 3-tridecylamino-1-propanesulfonic acid, 3-tetradecylamino-1-propanesulfonic acid, 3-hexadecylamino-1-propanesulfonic acid, and 3-octadecylamino-1-propanesulfonic acid.
- 5
93. The method of claim 89, wherein said A_t is selected from the group consisting of 1-phenyl-2,3-dimethyl-4-methylamino-pyrazolon-5-N-methylsulfonic acid; 3-dimethylamino-1-propanesulfonic acid, 3-[2-(1,2,3,4-tetrahydroisoquinoliny)]-1-propanesulfonic acid, 3-[2-(6,7-dimethoxy-1,2,3,4-tetrahydroisoquinoliny)]-1-propanesulfonic acid, 3-[1-(1,2,3,4-tetrahydroquinoliny)]-1-propanesulfonic acid, 2-(3-sulfopropyl)-1,2,3,4-tetrahydro-9H-pyrido[3,4-b]indole, 3-(1-indoliny)-1-propanesulfonic acid, 3-[2-(6-methoxy-1,2,3,4-tetrahydroisoquinoliny)]-1-propanesulfonic acid, 3-(2-isoindoliny)-1-propanesulfonic acid, 3-[2-(1,2,3,4,5,6,7,8-octahydroisoquinoliny)]-1-propanesulfonic acid, and 4-[2-(1,2,3,4-tetrahydroisoquinoliny)]-1-butanefulfonic acid .
- 10
- 15
94. The method of claim 63, where said A_t is selected from the group consisting of 4-phenyl-1-(3'-sulfopropyl)-1,2,3,6-tetrahydropyridine; 2-(3-sulfobutyl)-1,2,3,4-tetrahydro-9H-pyrido [3,4-b] indole; 3-(4-benzyl-1-piperidiny)-1-propanesulfonic acid, 3-sulfonylmethylphenylalanine, 4-(1-piperidiny)-1-butanefulfonic acid, cyclohexylsulfamic acid; 1-(3-sulfopropyl)-(S)-nicotinium hydroxide inner salt, 3-[1-(1,2,3,6-tetrahydropyridyl)]-1-propanesulfonic acid, 3-sulfomethyl-D,L-phenylalanine and 3-sulfomethyl-L-phenylalanine.
- 20
95. A method for diagnostic medical imaging of an amyloid-associated disease in a patient, comprising administering to a patient a pharmaceutical composition comprising an amyloid-targeting imaging agent of the formula
- 25



- 30 as defined in claim 63 and then imaging the amyloid-targeting imaging agent in said patient..

96. The method of claim 95, wherein A_{lab} of said pharmaceutical composition is a radiopharmaceutical.
97. The method of claim 95, wherein A_{lab} of said pharmaceutical composition is a metal chelate.
- 5 98. The method of claim 95, wherein A_{lab} of said pharmaceutical composition is a metal chelate and said imaging step is magnetic resonance imaging or radionuclide imaging.
99. The method of claim 97, wherein said metal chelate is gadolinium-DTPA, gadolinium-DOTA, or gadolinium-DO3A.
100. The method of claim 97, wherein said metal chelate is a chelate of ^{99m}Tc or ¹¹¹In.
- 10 101. The method of claim 63, wherein said imaging step is ultrasound imaging.
102. The amyloid-targeting imaging agent of claim 1, where A_i is 2-(3-sulfopropyl)-7-amino-1,2,3,4-tetrahydroisoquinoline, 3-[2-(5-amino-1,2,3,4-tetrahydroisoquinolyl)]-1-propane sulfonic acid, 2-(3-sulfopropyl)-6-amino-1,2,3,4-tetrahydro-9H-pyrido[3,4b]indole, 2-(4-sulfobutyl)-6-amino-1,2,3,4-tetrahydro-9H-pyrido[3,4b]indole, 1,6-hexanedisulfonate, 3-
15 hydroxypropylsulfamic acid, 4-(1-piperidinyl)-1-butanedisulfonic acid, 1,4-piperazine bis(propanedisulfonic acid), 3-[1-(1,2,3,6-tetrahydropyridinyl)]-1-propanedisulfonic acid, Thiazole yellow G, alpha-N-(3-sulfopropyl)-L-lysine, 3-(6-hydroxy-1-hexyl)amino-1-propane sulfonic acid, 3-(1-hydroxymethyl-1-cyclopentyl)amino-1-propane sulfonic acid, or methyl 2-(2-carboxyethyl)-1,2,3,4-tetrahydroisoquinoline hydrochloride, 2-(3-sulfopropyl)-7-amino-
20 1,2,3,4-tetrahydroisoquinoline, 3-[2-(5-amino-1,2,3,4-tetrahydroisoquinolyl)]-1-propane sulfonic acid, 2-(3-sulfopropyl)-6-amino-1,2,3,4-tetrahydro-9H-pyrido[3,4b]indole, or 2-(4-sulfobutyl)-6-amino-1,2,3,4-tetrahydro-9H-pyrido[3,4b]indole.